



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The greatest common divisor of these values is $\frac{1\frac{2}{3}50}{8}$.

$\therefore \$\frac{1\frac{2}{3}50}{8} = \$32\frac{2}{3}$ is the highest par value of each share in order that each may have whole shares and maintain his present interest.

A will then have 85 shares, B, C, D, E and E each 61 shares. Total number of shares in corporation, 624.

322. Proposed by THEODORE L. DeLAND, Treasury Department, Washington, D. C.

Take six consecutive prime numbers, as 53, 59, 61, 67, 71, and 73, and find the least whole number such that if it be divided by 59 the remainder will be 53, if it be divided by 67 the remainder will be 61, and if it be divided by 73 the remainder will be 71, and show that this least whole number and the succeeding consecutive whole numbers that will fulfill this condition as to divisions and remainders are in arithmetical progression; and also show whether or not this is a general law for n consecutive prime numbers; and if there be such a general law whether or not that general law will lead to a general law for the finding of prime numbers.

Solution by J. SCHEFFER, A. M., Kee Mar College, Hagerstown, Md.

Let x = the number. Then $\frac{x-53}{59}$, $\frac{x-61}{67}$, $\frac{x-71}{73}$ = whole number.

Proceeding in the usual way, we get $x = 27665 - 288569n$.

Therefore 27665 is the smallest whole number satisfying the condition. The succeeding numbers we get by putting $n = -1, -2, -3, \dots$ which, of course, form an arithmetical progression the common difference of which is 289569. This will always be the case for any number of consecutive prime numbers, though it will not lead to a general law for the finding of prime numbers.

GEOMETRY.

347. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

ABC is a triangle, and D, E, F are the mid points of the arcs of its nine-point circle cut off by BC, CA, AB , respectively. The inscribed circle touches these sides at X, Y, Z . Are the lines DX, EY, FZ concurrent? A purely geometrical discussion required.

No solution of this problem has been received.

348. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

Two parabolas and a rectangular hyperbola circumscribe a given quadrilateral. Find a relation between the squares of the latera recta of the parabolas and the squares of the perpendiculars from the center of the hyperbola to the axes of the parabolas.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

(1) $\dots m(x^2 + y^2) + 2nxy + 2px + 2qy + r = 0$ is the equation of the rectangular hyperbola;

(2) $\dots (ax + by)^2 + 2gx + 2fy + c = 0$ is the equation of one parabola, and

(3) $\dots (Ax + By)^2 + 2Gx + 2Fy + C = 0$ is the equation of the other parabola